

64455

Oriented, Glass-coated Impact Melt Rock

56.7 grams



Figure 1: Original PET photo of dusty glass egg 64455 in its upside down position (resting on zap pits). Scale is in cm. NASA # S72-40132. (compare with figure 11)

### **Introduction**

Sample 64455 has proven to be a nearly perfect sample for studies of the effects of cosmic ray, solar cosmic ray and micrometeorite bombardment of the lunar surface over the last 2 My, because it apparently maintained its orientation since it landed and been continuously exposed to the space environment for that time.

64455 is an egg-shaped object about 5 cm long and 3 cm across (figure 1) almost completely covered with thick black glass. The inside “yoke” is a fragment of basaltic melt rock, which is exposed in one quadrant on the top where the glass has broken off. The top surface is almost completely covered with micrometeorite craters, while the bottom surface has none, indicating that the sample has maintained its orientation since it landed on the lunar surface (figure 2).

### **Petrography**

Sample 64455 is a basaltic impact melt with very thick glass coat (Ryder and Norman 1980). Grieve and Plant (1973) describe four texture zones (a) interior fractured highland basalt, (b) basalt with interstitial partial melt, (c) a thin crust of brown devitrified glass, and (d) an outer coating of fresh glass (figure 3). Vaniman and Papike (1980) group the crystalline highland basalt interior fragment with the Low-K Fra Mauro suite and give the mineralogical mode as 59.9 vol. % plagioclase, 32.7 % pyroxene, 6.5 % olivine (with trace metal, troilite, schreibersite and “rust”).

### **Mineralogy**

The composition of pyroxene, olivine and plagioclase of the interior rock fragment are presented in Vaniman and Papike (1980) and Heiken et al. (1991) (figures 4 and 5). The composition of schreibersite and metal




Figure 2: Enlargement of portion of top surface (B1) of 64455 showing high density of micrometeorite "zap" pits and approximate position of saw cuts. Photo # S73-22656. Scale in mm is shown.




Figure 3: Photomicrograph of thin section of interior basalt fragment in 64455 (from Grieve and Plant 1973).






Figure 4: Pyroxene and olivine composition of interior "highland basalt" rock inclusion in 64455. Data replotted from Vaniman and Papike (1980) with apologies.




Figure 5: Plagioclase composition in basalt fragment in 64455 (from Vaniman and Papike, 1980).

grains are given in Grieve and Plant (1973), who also mention several metal grains showing alteration to a "rusty" component.

## Chemistry

Haskin et al. (1973) determined the major and trace element chemistry of the glass and interior rock fragment (table 1 and figure 6) and note that the glass can not simply be derived from the included rock fragment. Ganapathy et al. (1974) determined the trace and volatile composition and note that the volatile-rich glass may be from South Ray Crater (confirmed by Hertogen et al. 1977). The composition of the glass splashed on the outside of the rock has also been studied by See et al. (1986) and Morris et al. (1986).

The interior rock fragment does not appear to be "pristine" (Ir = 2.25 ppb).

## Radiogenic age dating

None

## Cosmogenic isotopes and exposure ages

The  $^{81}\text{Kr}$  exposure age of 64455 is 2.01 m.y. and the rock is associated with the South Ray cratering event




Figure 6: Normalized rare earth element plot for basaltic interior fragment and glass coating of 64455 (data from Haskin et al. 1973).

(Marti 1975, as reported by Arnold et al. 1993). Bogard and Gibson (1975) reported a  $^{21}\text{Ne}$  cosmic ray exposure age of 1.2 m.y. and  $^{36}\text{Ar}$  age of 1.8 m.y.

Nishiizumi et al. (1995) determined the  $^{10}\text{Be}$ ,  $^{26}\text{Al}$  (figure 7) and  $^{36}\text{Cl}$  activity by accelerator mass spectrometry of 19 sub-samples along a depth profile in 64455,82 (slab B). The distribution of microcraters and the  $^{10}\text{Be}$  activity are consistent with exposure of 64455 to cosmic rays for only 2 m.y. (Nishiizumi et al. 1995).

Arnold et al. (1993) also determined the  $^{10}\text{Be}$ ,  $^{26}\text{Al}$  and  $^{36}\text{Cl}$  as a function of depth below the surface of the protected side of 64455,82, by grinding away (peeling) thin surfaces with a drill.

## Other Studies

Neukum et al. (1973) studied the distribution of micrometeorite craters (figure 8). Blanford et al. (1974, 1975) studied the nuclear tracks (figure 9) and present a figure illustrating the lunar orientation (figure 10).

Leich et al. (1973) determined the depth distribution of H and F in exterior chips. Goldberg et al. (1976) attempted to determine solar wind implanted carbon.

## Processing

Two slabs have been cut from top to bottom of 64455 (figure 11). Slab A was prepared in 1972 (figure 12). Slab B was cut in 1992 (figure 13). Figure 14 illustrates the saw cuts on the T1 surface.




Figure 7: Depth profile for  $^{26}\text{Al}$  activity in 64455 (this is figure 2 from Nishiizumi et al. 1995).




Figure 8: Closeup photo of zap pits illustrating glass-lined pits with surrounding spall zones. Luckily, none of these impacts was large enough to disturb the orientation, although one may have been big enough to break the glass coating, exposing the interior rock sample. The uppermost surface was saturated with small craters (Neukum et al 1973). A few microcraters can also be seen on the exposed rock surface.




Figure 9: Cosmic ray particle tracks as a function of depth in 64455 (Blanford et al. 1975).





Fig. 1. A sketch of lunar sample 64455 in its supposed lunar orientation determined from its distribution of microcraters. Daughter samples 64455,14 and 64455,16 are shown as determined from cutting photographs taken by the curatorial staff. Dotted lines represent radii along which track density profiles were measured.

Figure 10: Orientation of 64455 from Blanford et al. (1975).

Note: The initial PET photography (with the sample in an upside down orientation, resting on the zap-pitted surface, figure 1) established the orientation cube with the B<sub>1</sub> surface being the zap-pitted surface. This has caused a lot of confusion.

## Sketch 64455



C Meyer 03

Figure 11: Schematic sketch of glass egg 64455 showing approximate positions of saw cuts to obtain two slabs for accurate depth profile studies of cosmogenic nuclides. Compare with figure 1.




Figure 12: Photo of E1 surface of slab B (64455,82), side next to slab A, showing vesicles in thick black glass and glass veins in rock. NASA # S91-36325




Figure 13: Photo of W1 surface of slab B (64455,82). from NASA # S91-36324 (orientation uncertain)

**Table 1. Chemical composition of 64455.**

reference weight	Haskin 73	Haskin 73 glass	Ganapathy 74 rock	Ganapathy 74 glass	Grieve and Plant 73 rock	Grieve and Plant 73 glass	Morris 86 glass		
SiO <sub>2</sub> %	48.5	44.6	(a)		47.17	45.2	(d)	44.76	(d)
TiO <sub>2</sub>	0.65	0.45	(a)		0.6	0.39	(d)	0.44	(d)
Al <sub>2</sub> O <sub>3</sub>	22.4	25.2	(a)		24.96	24.75	(d)	25.84	(d)
FeO	5.47	6.15	(a)		5.28	6.4	(d)	5.6	(d)
MnO	0.074	0.081	(b)		0.06	0.04	(d)	0.06	(d)
MgO	9.29	7.76	(a)		7.76	8.34	(d)	8.07	(d)
CaO	13.4	14.5	(a)		13.54	14.44	(d)	14.31	(d)
Na <sub>2</sub> O	0.57	0.37	(b)		0.28	0.44	(d)	0.36	(d)
K <sub>2</sub> O	0.245	0.106	(a)		0.21	0.05	(d)	0.13	(d)
P <sub>2</sub> O <sub>5</sub>									
S %									
sum									
Sc ppm	7.8	7	(b)					6.88	(b)
V									
Cr	1110	1000	(b)					899	(b)
Co	31.1	48.2	(b)					59	(b)
Ni	540	760	(b)	80	905	(c )		927	(b)
Cu									
Zn	4			2.2	2.4	(c )			
Ga	3.05	1.72	(b)						
Ge				62	500	(c )			
As									
Se				190	390	(c )			
Rb	6	3.1	(b)	6.6	3.9	(c )			
Sr									
Y									
Zr									
Nb									
Mo									
Ru									
Rh									
Pd ppb									
Ag ppb				1.2	1.6	(c )			
Cd ppb				5.3	5.2	(c )			
In ppb									
Sn ppb									
Sb ppb				0.45	3.6	(c )			
Te ppb				12.8	38	(c )			
Cs ppm	0.27	0.1	(b)	0.28	0.144	(c )			
Ba								122	(b)
La	21.1	12.6	(b)					12.09	(b)
Ce	56	32.1	(b)					35.1	(b)
Pr									
Nd	34	21	(b)						
Sm	9.8	5.9	(b)					5.88	(b)
Eu	1.23	1.11	(b)					1.14	(b)
Gd		7.4	(b)						
Tb	1.97	1.14	(b)					1.22	(b)
Dy	12.2	7.5	(b)						
Ho	2.7	1.6	(b)						
Er	6.7	4.5	(b)						
Tm									
Yb	6.6	3.92	(b)					3.76	(b)
Lu	0.96	0.56	(b)					0.57	(b)
Hf	7.8	4.3	(b)					4.13	(b)
Ta								0.51	(b)
W ppb									
Re ppb				0.284	4.11	(c )			
Os ppb									
Ir ppb				2.25	40.6	(c )			
Pt ppb									
Au ppb									
Th ppm								2.91	(b)
U ppm				1.43	0.86	(c )		0.54	(b)
technique (a) AA, (b) INAA, (c ) RNAA, (d) elec. Probe (broad beam)									





*Figure 14: Photo of the T1 surface (lunar bottom) of 64455 after cutting first slab A. The approximate location of the second slab is indicated by the line. The scale is in mm and the cube is 1 cm. There are no zap pits on this surface as it was below the soil line. Photo # S72-53870.*

NASA Photo #s	
S72-40130-40135	PET dusty
S72-43250-43266	PET B&W
S72-48510-48511	color close-ups
S72-53857	
S72-53870-53872	group photos
S73-22655-22659	
S89-39846-39849	
S91-36321-36325	saw cuts
S93-45928	
S93-45939-45942	